

Senegal Parrot

Poicephalus senegalus

Also known as yellow bellied Senegal parrot

Senegal parrots are small stocky parrots, one of the small *Poicephalus*. They are colorful parrots with a gray head, bright green upper parts, bright yellow orange abdomen and breast and green lower neck with a v shaped green patch extending down the chest.

Senegal parrots have a large range in western Africa from Guinea and Cameroon to Chad. They inhabit lowland forests from dry savannah to mosaic forests. Diet of wild birds is probably fruits, nuts and seeds. Wild birds are generally found in pairs or small groups and are shy and wary.

P. s. senegalus More eastern part of range

P. s. versteri – Ghana and Ivory Coast – Has more orange-red lower breast and abdomen and darker green uppers.

Length 9-9 ½ inches

Weight 125-170 grams. Average male weight – 156 grams. Average female weight 136 grams.

Life Span – possibly up to approximately 30 years but more likely approximately 15-20 years. Juvenile birds have dark gray eyes that become yellow as adults and many have more extensive green on the chest and abdomen.

Age of sexual maturity is 2-4 years.

Personality -. Senegals are playful outgoing birds that love attention. They can be affectionate but are not generally demanding of attention. They tend to become more independent as they reach sexual maturity and adult males may become aggressive during breeding season. While they are not great talkers, they have some limited mimicking ability.

Young senegals adapt readily to new surroundings and should be well adapted to many novel experiences at a young age. Adult birds are less adaptable to unfamiliar environments, dietary changes.

Activities - Senegals are very playful and energetic. Environmental enrichment is important. They should always be provided with toys, wooden blocks that can be chewed, and branches from non-toxic trees. In order to ensure safety, companion birds should not be allowed unsupervised freedom in the home as they often encounter toxins or dangerous items. Young birds should be socialized to many people and exposed to a variety of situations such as new cages, toys, visits to the veterinarian, handling by friends, wing and nail clips, etc to avoid fear of novel situations.

Dietary needs - African grays and to a lesser extent *Poicephalus* appear to utilize calcium differently than other psittacine species. Birds kept as indoor pets especially tend to develop signs of calcium deficiency that can be a serious health threat. Natural or full spectrum light may also be helpful. African parrots should be fed a formulated (pelleted or extruded diet) as a basis for good nutrition. Kaytee Exact is a complete and balanced diet for African parrots. Senegals should be fed approximately 1-2 heaping tablespoons of Kaytee Exact daily. Alternatively they can be fed Kaytee Exact small. Many birds will waste less food if fed smaller sized pellets. The diet should be supplemented with approximately the same quantity of fresh fruits and vegetables daily to add variety and psychological enrichment. Treats maybe given in small amounts especially as rewards for good behavior. Fresh clean water must be provided every day. Vitamin supplements are not needed for birds that are eating a formulated diet.

For birds fed a seed diet, vitamin supplementation is necessary. Vitaminized seeds have vitamins added to the shells that are discarded by the bird when it eats. Preferably vitamins should be added to soft foods rather than water as vitamins and the accompanying sweeteners promote bacterial growth in water.

Poicephalus can be very difficult to handrear from the egg and ideally should be fed by the parents for 1-2 weeks. If rearing from the egg they need to be fed very often, preferably every 1 ½ to 2 hours. Kaytee Exact Macaw Handrearing formula is a suitable hand-rearing diet. Senegals wean or fledge at 7-9 weeks.

Grooming - Routine bathing or showering is vital to maintaining good plumage and skin condition. Birds can be misted and allowed to dry in a warm room or in the sun, or dried with a blow drier. Care should be taken not to clip the wing feathers excessively as heavy bodied birds may fall and injure themselves. Clip only enough so the bird will glide to the floor.

Identification - All companion and breeding birds should be individually identified to assist in recovery if lost and assist in maintenance of medical and genealogy records. Many breeders apply closed legs bands when chicks are young. While they present a slight risk of entrapment closed bands are preferable to no identification, especially for breeding birds. Microchips, which can be implanted into the muscle or under the skin, are a reliable means of identification but require electronic readers to verify identification. Tattoos may be used but often fade or become illegible with time. Footprints may have some application in identification.

Sexing - Senegal parrots show some slight sexual dimorphism (visual difference between the species) however for breeding birds, endoscopic examination or laboratory sexing techniques are needed for accurate sex determination. Males tend to have solid yellow under-tail coverts. Females tend to have more green in the under-tail coverts. For photos visit the web page of the African Parrot Society.

Housing - African parrots are very active and should be provided with as large a cage as possible. The cage should have two perches so the birds can move between them. Toy

and activities should be provided. Ideally pet birds should have a cage outdoors to allow exposure to sunlight and fresh air in good weather.

Breeding - Senegal parrots breed well in captivity. Some prolific birds will breed year round but most breed in the winter and early spring. Clutch size is usually 3-4 eggs.

Nest Box - Senegals will use a vertical 10" x 10" x 12" or an L shaped box.

Cage size - Cage size should be at least 4' x 4' x 4' or 3' x 3' x 6'.

Common Diseases

- Respiratory Diseases- Aspergillosis
- Bacterial, viral, Fungal Diseases
- Calcium deficiency disorder
- Toxicities
- Chlamydiosis (Psittacosis)
- Psittacine Beak and Feather Disease
- Feather picking

Many common health problems can be prevented by good diet, nutrition and routine health care. Routine veterinary examination (annually) can help you to keep your pet in excellent health and enhance your relationship with your bird.

Conservation Status – Common , Stable – Senegal are listed on Appendix II of CITES (Convention on Trade in Endangered Species of Fauna and Flora) because of the listing of almost all parrots. They are common in the market place and increasing in popularity as pets.